Southeast Tennessee HIV Community Planning Region, 2022




65 persons were newly diagnosed with HIV









25% received a late diagnosis

→ Persons diagnosed with HIV Stage 3 (AIDS) within 12 months of a new HIV diagnosis are considered a **late diagnosis**, indicating delayed awareness of HIV status and missed opportunities for prevention efforts.

Most affected subpopulations by number of persons newly diagnosed with HIV

Non-Hispanc black men, MMS				13
Non-Hispanic white men, MMS 9				
Non-Hispanis black men, HET 7				
6 Non-Hispanic white men, MMS+IDU				
5 Hispanic men, MMS				
4	Non-Hispanic white women, IDU			
4	Non-Hispanic white men, IDU			
MMS: male-to-male sexual contact; HET: heterosexual contact; IDU: injection drug use				

68% were linked to HIV care within 30 days of diagnosis

91% were linked within 365 days

1,726 persons were living with diagnosed HIV

82%

received any HIV medical care

→ One lab result reported to TDH

61%

were retained in HIV medical care

→ Two or more lab results, at least 90 days apart, reported to TDH

62% were virally suppressed

→ When HIV medication reduces the amount of HIV virus in a person's body to a low enough level (<200 c/mL), they are considered **virally suppressed** and will not transmit HIV through sexual contact.





For more HIV surveillance data visit: tn.gov/health/HIVData